

The scope was inserted up to the cecum with visualization of the appendiceal orifice and ileocecal valve. The terminal ileum was intubated, which showed normal mucosa. On withdrawal, a polypoid mass with friable mucosa that obstructed approximately 60-70% of the lumen was noted at the mid-transverse colon, from which multiple biopsies were taken for histopathologic examination. The rest of the visualized colonic mucosa was pinkish, smooth and with regular vascular pattern. The hemorrhoidal vessels above the dentate line were slightly congested.

The patient tolerated the procedure well, which was completed in 15 minutes with withdrawal time of 10 minutes, and with no untoward reactions noted.

Medications: Midazolam 2 mg IV; Ketamine 50 mg IV
Complication/s: Minimal bleeding from biopsy sites, which resolved spontaneously.

Post-endoscopy Diagnosis:

POLYPOID MASS, TRANSVERSE COLON, PROBABLY MALIGNANT, S/P BIOPSY
INTERNAL HEMORRHOIDS, GRADE 1

Colorectal Surgery of my mothyer

Our goal is to raise funds for my mother’s urgent colorectal surgery and hospital expenses

Donate on the fundraising campaign

Scan and Donate

Point your phone's camera at the QR code.

Or go to

steunactie.be/en/-54363

Fundraiser organised by

Marifi Guan